

<https://helda.helsinki.fi>

Aeolian dust supply from the Yellow River floodplain to the Pleistocene loess deposits of the Mangshan Plateau, central China : Evidence from zircon U-Pb age spectra

Shang, Yuan

2018-02-15

Shang , Y , Prins , M A , Beets , C J , Kaakinen , A , Lahaye , Y , Dijkstra , N , Rits , D S , Wang , B , Zheng , H & van Balen , R T 2018 , ' Aeolian dust supply from the Yellow River floodplain to the Pleistocene loess deposits of the Mangshan Plateau, central China : Evidence from zircon U-Pb age spectra ' , Quaternary Science Reviews , vol. 182 , pp. 131-143 . <https://doi.org/10.1016/j.quascirev.2018.01.001>

<http://hdl.handle.net/10138/309290>

<https://doi.org/10.1016/j.quascirev.2018.01.001>

cc_by_nc_nd

acceptedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.

Please cite the original version.

Aeolian dust supply from the Yellow River floodplain to the Pleistocene loess deposits of the Mangshan Plateau, central China: Evidence from zircon U-Pb age spectra

Yuan Shang^{a,b,*}, Maarten A. Prins^b, Christiaan J. Beets^b, Anu Kaakinen^a, Yann Lahaye^c, Noortje Dijkstra^{b,1}, Daniël S. Rits^b, Bin Wang^d, Hongbo Zheng^e, Ronald T. van Balen^{b,f}

^a *Department of Geosciences and Geography, P.O. Box 64, 00014, University of Helsinki, Finland*

^b *Department of Earth Sciences, Faculty of Science, Vrije Universiteit Amsterdam, De Boelelaan 1085, 1081 HV Amsterdam, The Netherlands*

^c *Geological Survey of Finland (GTK), Betonimiehenkuja 4, 02150 Espoo, Finland*

^d *School of Geography and Tourism, Shaanxi Normal University, 710119, Xi'an, China*

^e *School of Resource, Environment and Earth Science, Yunnan University, 650500, Kunming, China*

^f *TNO-Geological Survey of the Netherlands, Princetonlaan 6, 3584 CB Utrecht, The Netherlands*

* Corresponding author: yuan.shang@helsinki.fi; y.shang@vu.nl

¹ Present address: Department of Geosciences, UiT the Arctic University of Norway in Tromsø, Dramsveien 201, 9037 Tromsø, Norway

Abstract

The thick loess-palaeosol sequences in the Mangshan Loess Plateau (MLP; central China) along the south bank of the lower reach of the Yellow River provide high-resolution records of Quaternary climate change. In addition, substantial increases in grain-size and accumulation rate have been inferred in the upper part of the loess sequence, above palaeosol layer S2. This study investigates the sources of the long-term dust supply to the MLP and explores the mechanism behind the sudden increase in sediment delivery and coarsening of the loess deposits since S2 (~240 ka) by using end member modelling of the loess grain-size dataset and single-grain zircon U-Pb dating. Our results indicate that the lower Yellow River floodplain, directly north of the MLP, served as a major dust supply for the plateau at least since the deposition of loess unit L9 and indirectly suggest that the integration of the Yellow River and the disappearance of the Sanmen palaeolake took place before L9 (~900 ka). The sudden change in sedimentology of the Mangshan sequence above palaeosol unit S2 may result from an increased fluvial sediment flux being transported to the lower reaches of the Yellow River

because of tectonic movements (initiated) in the Weihe Basin around 240 ka. Furthermore, sediment coarsening can be explained by the gradual southward migration of the lower Yellow River floodplain towards the MLP since the deposition of palaeosol S2. The migration is evidenced by the formation of an impressive scarp, and is likely caused by tectonic tilting of the floodplain area.

Key Words: Provenance analysis; Sanmen Gorge; Mangshan Plateau; end member modelling; grain-size

1. Introduction

Quaternary loess-palaeosol sequences are widespread in the Chinese Loess Plateau (CLP) of northern China. The sedimentological characteristics of these wind-blown deposits provide a powerful tool for the reconstruction of the past atmospheric circulation pattern and climate change (Liu, 1985; Kukla, 1987; An et al., 2001; Porter, 2001). A SE-ward decreasing trend in grain size, unit thickness and inferred sedimentation rate is recognised in the CLP loess sequences (Liu, 1985; Pye, 1995; Liu and Ding, 1998; Ding et al., 2002; Nugteren and Vandenberghe, 2004; Prins and Vriend, 2007). Two factors mainly control grain-size variations within the CLP, namely the strength of transporting winds and the locations of the source areas. Therefore, the sedimentary characteristics of the wind-blown deposits allow us to assemble information concerning past atmospheric circulation patterns and the distance to the source area. Previous studies suggested that the deserts and arid lands north and northwest of the CLP are the main source area of the CLP loess deposits (Liu, 1985; Derbyshire et al., 1998; Lu and Sun, 2000; Sun, 2002; Nugteren and Vandenberghe, 2004; Ding et al., 2005). However, recent evidence based on single grain zircon U-Pb dating found a genetic link between the Yellow River and loess sediments. These results emphasise the important contribution of reworked fluvial detritus delivered by the Yellow River from the NE Tibetan Plateau to the CLP (Stevens et al., 2013; Bird et al., 2015; Nie et al., 2015; Licht et al., 2016; Zhang et al., 2016).

The Yellow River is the second longest river of China, with a total length of 5500 km. It originates in the northeast Tibetan Plateau and runs eastwards around the Ordos block and the North China Plain before discharging into the Bohai Sea (Fig.1). The river course has traditionally been divided into upper, middle and lower reaches, with Hekou town in Inner Mongolia marking the upper/middle reach boundary, and Mengjin

60 of Henan province the middle/lower reach boundary of the river (Fig. 1). As the most sediment-laden river in
61 the world, the Yellow River delivered more than one billion tonnes sediments each year to the sea between
62 1951 and 1979 (Wang et al., 2015). Its sediment load increases most markedly in the middle reach, near the
63 CLP and gradually declines downstream.

64 Geological surveying has identified a palaeolake in the Weihe Basin, with its eastern boundary at the
65 Sanmen Gorge (AFSOM, 1988; Jiang et al., 2007). It has been suggested that the Sanmen palaeolake formed
66 ~5 Ma (Wang, 2002) and was drained when the Yellow River started to cut the Sanmen Gorge, leading to the
67 formation of its modern course (Jiang et al., 2007; Zheng et al., 2007; Wang et al., 2013). The proposed timing
68 of the incision of the gorge varies from late Miocene to Pleistocene (Lin et al., 2001; Wang, 2002; Jiang et al.,
69 2007; Wang et al., 2013; Kong et al., 2014; Hu et al., 2017), and Rits et al. (2017) concluded that the Sanmen
70 Lake did not exist during the last ~1 Ma. The cutting of the Sanmen Gorge allowed the release and transport
71 of substantial volumes of reworked loess towards the lower reaches of the river. As a result, large fluvial fan
72 systems were created east of the Sanmen Gorge (Huang et al., 2009). Studying the evolution of the Yellow
73 River is crucial for understanding the “source-to sink” process, influenced by tectonic activity and climate
74 change at both a regional and global scale. However, the geological history of the Yellow River remains a
75 topic of debate and active investigation, particularly for the development of its drainage system in its middle
76 and lower reaches (Lin et al., 2001; Wang, 2002; Jiang et al., 2007; Pan et al., 2011; Wang et al., 2013; Kong
77 et al., 2014; Hu et al., 2017; Li et al., 2017).

78 Thick, continuous loess-palaeosol records of exclusively aeolian origin have been found on the Mangshan
79 Loess Plateau (MLP) (Wu et al., 1999; Jiang et al., 2007; Zheng et al., 2007; Qiu and Zhou, 2015) which is
80 located along the lower reach of the Yellow River (Fig. 1). Here the loess deposits above palaeosol S2
81 (equivalent to MIS 7, ca. 225 ka) are significantly thicker than loess deposits on the CLP. For instance, the
82 deposits of L1-S1-L2 of Xifeng, Luochuan and Lantian are between 10 and 20 m thick, whilst those at
83 Mangshan are ~86 m thick. Consequently, sedimentation rates and mass accumulation rates are significantly
84 higher at the MLP than in the central CLP (Prins et al., 2009). In addition, the upper part of the Mangshan
85 record displays extremely high sedimentation rates and a coarser composition compared to its lower part,
86 suggesting a dust supply from a more proximal source, i.e. the lower Yellow River floodplain (Wu et al., 1999;
87 Jiang et al., 2007; Zheng et al., 2007; Prins et al., 2009). Study of the provenance and sedimentological

variations of the Mangshan loess records may therefore assist in the understanding of the mechanisms associated with dust supply from the Yellow River to the MLP and the drainage system development in the river's middle and lower reaches.

In this paper, the first record of single grain zircon U-Pb chronology of the Mangshan loess-palaeosol sequence is presented (Figs. 2 and 5, from the Holocene soil S0 through L6 and L9) and is compared with the U-Pb age signature of possible sources, including the upper, middle and lower reaches of the Yellow River deposits, the CLP loess, as well as river sediments from the nearby Taihang Mountains and Qinling Mountains. By combining the provenance age distributions with a mixing model of the grain-size distribution data (Figs. 2 and 3) and a dust flux model (Fig. 4) of the Mangshan sequence(s), it is intended to 1) characterise the contribution of subpopulations of the sediments and their corresponding transporting processes based on the grain-size dataset, 2) investigate the provenance signal of the Mangshan loess, comparing it to potential source areas (Fig. 6) and quantify their contributions (Fig. 7), and 3) discuss the mechanisms controlling sedimentology and provenance variations of the Mangshan dust during the Pleistocene and Holocene. The results also have implications for the evolution of the Yellow River system and the age of the Sanmen palaeoake.

2. Material and methods

2.1 Sites, samples and sediment analyses

The MLP lies 25 km west of Zhengzhou on the south bank of the Yellow River (Fig. 1). The loess plateau is about 18 km in length (W-E) and 5 km in width (N-S), with its highest point reaching approximately 150 m above the Yellow River floodplain (Fig. S1). The deposits consist of a number of loess and palaeosol alternations with a total thickness of ~170 m (Jiang et al., 1998; Ji et al., 2004; Jiang et al., 2004; Jiang et al., 2007; Zheng et al., 2007). Zheng et al. (2007) showed that the upper part of the sequence (0-97 m, including palaeosols S0, S1 and S2, and loess units L1 and L2) experienced extremely high sedimentation rates (on average 40 cm/kyr) in comparison with the lower part of the profile (97- 165 m; palaeosols S3-S10, loess units L3-L10) (see also Fig. S1).

116 Here the grain-size, magnetic susceptibility, carbonate content and organic matter records for the loess-
117 palaeosol sequences are presented, based on samples from two localities (Fig. 1 and Figs. S1, S2 and S4). The
118 northern ('proximal') loess section, here referred to as MS2006 (34°57.5' N, 113°22.2' E) (Fig. 1c), is exposed
119 on the northern slope of the plateau, where the Yellow River and local gullies cut through the loess forming a
120 valley with steep cliffs. Section MS2006 is ~130 m thick (Fig. S2) and is a composite of record X (0-59 m;
121 Prins et al., 2009) and record Z' (36-130 m; Zheng et al., 2007, see also Prins et al., 2009), analysed at a 10-
122 cm (X: 0-59 m) and 20-cm resolution (Z': 59-130 m). The grain-size, organic matter and carbonate profiles of
123 records X and record Z' for the overlapping interval 36-59 m indicate the similarities between records X and
124 Z' (Fig. S3). The uppermost S0-L1-S1 loess-palaeosol complex has also been sampled at a more southern
125 location, about 2.0 to 2.7 km south of the MS2006 section (Fig. 1c and Figs. S1 and S4) to study the impact of
126 increasing distance with respect to the Yellow River. The two sections MS2008W (34°56.4' N, 113°22.2' E)
127 and MS2008E (34°56.1' N, 113°22.4' E), ~34 and ~14 m thick, respectively, have been sampled in 26 partly-
128 overlapping vertical trenches at a 10-cm resolution.

129

130 **2.2 Magnetic susceptibility, carbonate and organic matter analysis**

131

132 The samples were dried in an oven (50 °C), lightly ground and aliquots of ~8 g were analysed using a
133 Bartington MS2 magnetic susceptibility meter at the School of Ocean and Earth Sciences, Tongji University
134 (Shanghai). The organic matter and carbonate content of aliquots of ~2 g of these samples were analysed by
135 thermo-gravimetric analysis (TGA), using a Leco TGA 601 at the Vrije Universiteit Amsterdam (VUA).
136 Derivative weight loss curves most clearly indicate the temperature intervals during which weight loss is most
137 apparent: replicate tests performed on the loess samples indicate that the temperature interval during which
138 organic matter and carbonate minerals disintegrate is 200-550 °C and 700-1000 °C, respectively.

139

140 **2.3 Grain-size analysis and end-member modelling**

141

142 The Mangshan loess samples for grain-size analysis were prepared following the methods described by
143 Konert and Vandenberghe (1997), with organic matter and carbonates being removed. All the measurements

were performed on a Fritsch Analysette 22 laser particle sizer at the VUA resulting in a grain-size distribution with 56 size classes in the size range 0.15-2000 μm .

A mixing model of the combined Mangshan grain-size distribution dataset ($n=1931$, from sections MS2006, MS2008W and 2008E) was constructed with the inversion algorithm for end-member modelling of compositional data EMMA (Weltje, 1997). EMMA is a non-parametric numerical-statistical technique and its advantage over the parametric curve fitting approaches, e.g. Weibull (Sun et al., 2002, 2004) or log-normal (Xiao et al., 2009, 2013) is that it does not require any prior knowledge about the grain size controlling processes (Weltje and Prins, 2007). This method has proven to be powerful in distinguishing aeolian from fluvial sediments in various marine settings (e.g. Prins and Weltje, 1999; Prins et al., 2000; Stuut et al., 2002, 2014; Deplazes et al., 2014) and in partitioning multiple transport/deposition processes of Quaternary loess from the CLP (Prins and Vriend, 2007; Prins et al., 2007; Vriend et al., 2011; Shang et al., in press). Details of the technical aspects of the end-member modelling algorithm are given in Weltje (1997), Prins and Weltje (1999) and Weltje and Prins (2003). Prins et al. (2009) already applied this approach to the grain-size data of the upper 60 m of the MS2006 section (Fig. 3d). In this study, a new mixing model expressing the loess samples from all three sections as mixtures of three end members has been produced.

The EMMA approach involves two modelling stages. In the first stage, the number of end members (EMs) is estimated on the basis of the mean and/or median coefficient of determination statistics (r^2 , Fig. 3a). The coefficients of determination represent the proportions of the variance of each variable (size class) that can be reproduced by the approximated data (Fig. 3b). In principal, the simplest model is chosen when the r^2 shows satisfactory goodness of fit (usually $r^2 > 0.8$, e.g. Fig. 3a). In the second modelling stage, the compositions of the end members are estimated (Fig. 3c) and the proportional contributions of the end members in the analysed samples are calculated (Fig. 2c).

166

167 **2.4 Age model and dust flux calculations**

168

Peterse et al. (2011) presented an age model for the upper 0-60 m (S0-L1-S1) of the sequence in section MS2006, and sections MS2008W and MS2008E based on the correlation of the typical loess proxy records, i.e. magnetic susceptibility, carbonate content and grain-size, with the $\text{U-}^{230}\text{Th}$ dated oxygen isotope records

from Dongge, Sanbao and Hulu caves in central China (Cheng et al., 2009; Wang et al., 2008). Following a similar approach, we extended the age model for the upper 130 m (S0-L6) of the loess-palaeosol sequence MS2006 (Fig. 2a, b, c) based on the correlation of loess proxy records with the newly published speleothem records extended to 640 kyr B.P. in central China (Fig. 2d; Cheng et al., 2016, and references therein). The inflection points in the loess proxy records have been used as tie points between the loess record and the MIS boundaries recognised in the stacked speleothem $\delta^{18}\text{O}$ record. The depth and ages of the 11 selected time control points are listed in Table S1. Simultaneously, the loess-palaeosol sequence was also visually correlated with the stacked benthic oxygen isotope record (Lisiecki and Raymo, 2005) assuming that the palaeosols layers (S0, S1 ... S5) correlate to interglacial Marine Isotope Stages (MIS1, 5...13-15) and the loess layers (L1, L2 ... L5) correlate to glacial Marine Isotope Stages (MIS 2-4, 6 ... 12) (Fig. 2e). Although a fully independent age control for the complete studied sequence is missing, a recent study by Qiu and Zhou (2015) provided OSL ages for the upper ~120 m of another section (covering the S0 to S5 interval) on the Mangshan Plateau. Their independent age model (based on the application of an elevated temperature post-IR IRSL ($p\text{IR}_{200}\text{IR}_{290}$) SAR procedure to polymineral fraction) is very similar to our findings presented in Figure 4 indicating that our age model suffices for the purpose of dust flux calculations.

Mass-accumulation rates (MAR, in $\text{g}/\text{cm}^2/\text{ka}$) of the well-constrained loess (L1, L2 ... L5) and palaeosol (S0, S1 ... S5) units of sections MS2006 and MS2008W were calculated according to Prins and Vriend, 2007:

$$(1) \quad MAR = SR \times BD$$

Where SR is the sediment accumulation rate (in cm/ky), and BD is the sediment dry-bulk density (in g/cm^3). SR values were calculated based on the age estimates (Table S1) and sediment thickness values for each of the loess and palaeosol intervals (Table S2). BD values of $1.48 \text{ g}/\text{cm}^3$ were used (cf. Kohfeld and Harrison, 2003).

Fractionated mass-accumulation rates (fluxes) for the modelled end members ($F_{\text{EM-x}}$, in $\text{g}/\text{cm}^2/\text{ka}$) were calculated according to Prins and Vriend (2007):

$$(2) \quad F_{\text{EM-x}} = MAR \times p_{\text{EM-x}}$$

Where $p_{\text{EM-x}}$ is the proportional contribution (dimensionless) of end member EM-x, and $\sum_{x=\text{EM-1}}^{\text{EM-3}} p_x = 1$.

197 In these calculations it was assumed that the loess-palaeosol samples are entirely composed of siliciclastics,
198 which is just a first-order approximation as the contribution of other (non-siliciclastic) sediment phases like
199 (detrital, pedogenic) carbonates and organic carbon has been ignored.

200 **2.5 Zircon samples and zircon U-Pb dating**

201 The zircon samples were collected from the MS2006 section, from loess units L1, L2, L3, L5, L6 and L9
202 (as a reference sample for the bottom loess unit, Fig. 2 and 5). Samples collected from potential source areas
203 in this study are shown in Figure 1b, including fluvial samples from the lower Yellow River, Yiluo River, Qin
204 River and the Yu River alluvial fan northeast to the MLP. The detailed location and description of zircon
205 samples including those from Kong et al. (2014), Bird et al. (2015) and Nie et al. (2015) are shown in Table
206 S3.

207 The zircon grains in the size range of 20 μm to 90 μm were extracted following the standard procedure of
208 heavy liquids and Frantz magnetic separation at the Mineral Separation Laboratory of VUA (Shang et al., 2016)
209 and randomly selected by hand-picking under an optical microscope and then mounted in epoxy resin and
210 sectioned approximately in half and polished. Back-scattered electron images (BSE) were prepared for the
211 zircons to target the spot sites. U–Pb dating analyses were performed using a Nu Plasma AttoM single collector
212 ICPMS connected to a Photon Machine Excite laser ablation system at the Geological Survey of Finland in
213 Espoo. Typical ablation conditions were: beam diameter: 20 μm , pulse frequency: 5 Hz, beam energy density:
214 2 J/cm². Raw data were corrected for the background, laser induced elemental fractionation, mass
215 discrimination and drift in ion counter gains and reduced to U–Pb isotope ratios by calibration to concordant
216 reference zircons, using the program Glitter (Van Achterbergh et al., 2001). Age related common lead (Stacey
217 and Kramers, 1975) correction was used when the analysis showed common lead contents significantly above
218 the detection limit (i.e., >50 cps). All the ages were calculated with 2 σ errors and without decay constants
219 errors. ²⁰⁶U/²³⁸Pb and ²⁰⁷Pb/²⁰⁶Pb ages were used for ages younger and older than 1 Ga, respectively. We used
220 a 10-20% discordance filter to the generated data.

221 **3. Results**

222 **3.1 Mangshan loess-palaeosol stratigraphy and sedimentology**

223 The carbonate, organic matter and median grain-size records of section MS2006 are shown in Figure S2.
224 On the basis of these parameters a clear distinction can be made between the loess and palaeosol layers.
225 Palaeosol layers S0 to S5 characteristically consist of fine-grained sediments, i.e. median grain-size values
226 below 25 μm , with low carbonate contents (1–2 %) and slightly, but consistently, elevated organic matter
227 contents (up to 1.5–2 %). In contrast, loess layers L1 to L6 consist of coarser sediments, with median grain-
228 size values up to 30–35 μm in the lower part of the sequence (L3–L6), and up to ~ 45 μm in loess horizons L2
229 and L1. The loess sediments typically show carbonate contents of $\sim 10\%$, and low organic matter values (~ 0.5 –
230 0.75 %). Distinct layers with high carbonate content ($>10\%$) – reflecting the presence of carbonate nodules –
231 occur near the base of palaeosol layers S1, S2, S3 and S4.

232 A detailed stratigraphic picture of the S0-L1-S1 sequence in section MS2006, and its correlation with
233 sections MS2008W and MS2008E, is shown in Figure S4. Overall, the three sections show relatively consistent
234 trends in the analysed proxies during the last glacial-interglacial cycle: glacial loess unit L1 is characterised
235 by relatively lower magnetic susceptibility of ~ 5 SI/g, higher carbonate content of $\sim 10\%$ and a coarser median
236 grain size (30–50 μm), while a higher magnetic susceptibility (5–10 SI/g), lower carbonate content ($<10\%$)
237 and finer median grain sizes (20–40 μm) characterise the interglacial palaeosol units S1 and S0. In addition,
238 higher-frequency and lower-amplitude patterns are visible in all three proxy records, within palaeosol unit S1
239 and especially in the loess unit L1, pointing to highly variable sedimentation and pedogenic processes
240 throughout the last glacial-interglacial cycle. The thickness of the loess unit L1 and palaeosol unit S1 decreases
241 from the northern MS2006 section to the southern MS2008W and MS2008E sections (Figs. S1 and S4). For
242 example, the thickness of L1 of the MS2006 section is about 42 m while in the MS2008W, it is around 22 m.

243 3.2 Mixing model

244 The ‘goodness-of-fit’ (r^2) statistics were computed for mixing models with 2–10 end members. The results
245 show that the loess sediments can be adequately described as mixtures of three end members (Figs. 3a and 3b).
246 The three-end-member model explains on average more than 86% of the observed variance in the grain-size
247 dataset (mean r^2 of 0.86 and a median r^2 of 0.92, Fig. 3b).

248 The end members are characterised by unimodal, fine skewed grain-size distributions (Fig. 3c) with modal
249 particle sizes close to 63 μm (EM-1), 37 μm (EM-2) and 26 μm (EM-3). The sand (>63 μm): silt (8–63 μm):

250 clay ($<8\ \mu\text{m}$, cf. Konert and Vandenberghe, 1997) ratio for EM1, EM2 and EM3 are 32:54:14, 15:76:9 and
251 2:55:43, respectively. The modelled end members presented here for the complete Mangshan dataset (Fig. 3c)
252 resemble those of the upper (S0-L1-S1) Mangshan sequence (Fig. 3d) presented by Prins et al. (2009) and the
253 CLP loess dataset (Fig. 3e) presented by Prins and Vriend (2007).

254 The proportional contribution of the end members with depth in the MS2006 section is compared to the
255 loess-palaeosol stratigraphy and median grain-size record in Figure 2. An evidently dramatic increase of the
256 sandy EM-1 in the loess units and palaeosol units above S2 is present. By contrast, the proportion of clayey
257 EM-3 decreased in the palaeosol units S0 and S1 compared to the lower units S2, S3, S4 and S5 (Fig. 2c).

258 A detailed picture of the S0-L1-S1 sequence in sections MS2006, MS2008W and MS2008E is depicted in
259 Figure S5. The proportion of the sandy EM-1 in MS2006 is significantly higher than in the southern two
260 sections MS2008W and MS2008E which are situated at a more distant location with respect to the Yellow
261 River floodplain (Fig. 1c and Fig. S1), whereas the proportions of the silty EM-2 increase from the northern
262 MS2006 section to MS2008-W and MS2008-E (Fig. S5). The clayey EM-3 does not show a clear spatial
263 increasing/decreasing trend. These trends suggest that the sandy and silty end-members of the upper three units
264 (S0-L1-S1) of the Mangshan loess were derived from a nearby source area, the lower Yellow River floodplain,
265 see below in the Discussion.

266 3.3 Dust flux model

267 Linear sediment accumulation rate (LSR), mass accumulation rates (MAR) and fractionated fluxes of the
268 three end-members ($F_{\text{EM-1}}$, $F_{\text{EM-2}}$ and $F_{\text{EM-3}}$) were calculated for the stratigraphic units of section MS2006 (S0-
269 L1-S1... L5-S5) and MS2008W (S0-L1-S1). Overall, a dramatic increase in the LSR is observed in the upper
270 part of MS 2006 section (above S2) (Fig. 4a). The average SR value for the upper part of MS2006 section is
271 36 cm/kyr while for the lower part (below S2) it is about 11 cm/ kyr. In more detail, the SR record shows an
272 increasing trend from the base (S5) to the top (L1), superimposed on a clear glacial-interglacial variability with
273 high LSR recorded during glacials (loess units) and low LSR during interglacials (palaeosol units).

274 Figure 4b shows that bulk MAR values range between ~ 4 and $\sim 88\ \text{g/cm}^2/\text{kyr}$ in section MS2006, with
275 minimum values in the S5 palaeosol unit and maximum values in the L1 loess unit. The MAR values in section
276 MS2008W varied between ~ 22 and $\sim 53\ \text{g/cm}^2/\text{kyr}$, with minimum values in the S1 palaeosol unit and

277 maximum values in the L1 loess unit. Similar to the SR record, the loess units display higher MAR than the
278 palaeosol units. However, there is a significant increase in MAR from S2 and above. Note that the MAR values
279 in palaeosol units S2 (25 g/cm²/kyr) and S1 (50 g/cm²/kyr) are even higher than the pure loess units L4 (9
280 g/cm²/kyr) and L5 (22 g/cm²/kyr). Spatially, the loess unit L1 and palaeosol unit S1 of the MS2006 section
281 also display overall higher MAR values compared to the units of MS2008W section, which is at a greater
282 distance from the Yellow River.

283 The summed fractionated fluxes of the coarse fraction ($F_{cf} = F_{EM-1} + F_{EM-2}$) are plotted against the bulk
284 MARs of the stratigraphic units in the sections MS2006 and MS2008W in Figure 4b. The coarse fraction and
285 MAR are positively related by the linear regression equation: $MAR = 1.10F_{cf} + 6.12$ ($r^2 = 0.98$). “6.12” reflects
286 the constant absolute contribution of EM-3 over time. The variations in relative EM-3 content are thus
287 dominantly caused by a variable input of end-members EM-1 and EM-2, with a negative relation between EM-
288 3 and MAR. Similar results have been described by Prins and Vriend (2007), Prins et al. (2007) and Vriend et
289 al. (2011) from the CLP.

290 **3.4 The zircon U-Pb age distributions**

291 **3.4.1 Mangshan loess-palaeosol sequence**

292 Figure 5 displays the zircon U-Pb age distributions for the Mangshan loess samples next to the loess-
293 palaeosol stratigraphy. All the loess samples are characterised by two dominant age populations, one at 200–
294 350 Ma (Permian–Triassic population) and the other at 350–550 Ma (Ordovician–Silurian population).
295 Additionally, there are several minor age peaks in the ranges of 0–100 Ma, 0.7–1 Ga, 1.5–2.0 Ga and at ca. 2.5
296 Ga. However, the relative abundance of the two dominant age peaks (200–350 Ma and 350–550 Ma) varies
297 from sample to sample. The abundance is almost equal for the samples from the loess units L1 (MS-L1-1 and
298 MS-L1-2) and L3 (MS-L3) whereas the age population at 200–350 Ma is more notable in sample MS-L2-1
299 from the upper part of loess unit L2, and the peak at 350–550 Ma is dominant in the samples from the lower
300 part of L2 (L2-2), the loess units L5 (MS-L5), L6 (MS-L6) and L9 (MS-L9).

301 In order to minimise uncertainty introduced by varying number of grains analysed for each sample ($n=213$ –
302 347), and to make the results more statistically meaningful, the zircon ages were grouped into an upper and a

303 lower Mangshan section. The comparison of the combined age spectra above S2 (n=1100) to those below S2
304 (n=903) shows that the age populations 200–350 Ma and 350–550 Ma are dominant in both datasets (Fig. 6a
305 and 6b), although, the <100 Ma and the 200–350 Ma age populations slightly increase (compared to the 350–
306 550 Ma population) in the upper part of the Mangshan sequence.

307 **3.4.2 Comparison of Mangshan loess with potential source areas**

308 Grain-size analyses and end-member modelling indicated that the nearby lower Yellow River floodplain
309 is the likely source of the sandy and silty loess components. The comparison of the zircon U-Pb age spectra of
310 the Mangshan loess with potential source areas (Fig. 6) confirm this and indicate that the overall pattern of
311 Mangshan zircon U-Pb ages (Figs. 6a and 6b) is not only comparable to that of the lower reaches of the Yellow
312 River as sampled at locations YR-20 and YR-33 (Fig. 1 b; Figs. 6k and 6l) but also to that of the loess deposits
313 from the CLP (Beiguoyuan and Lingtai, Fig. 1; Fig. 6c), and even the upper reaches of the Yellow River
314 (Fig. 6m). All these deposits exhibit two distinct Palaeozoic age populations at 200–350 Ma and 350–550 Ma
315 (Figs. 6k, 6l and 6o). Additionally, the upper part of Mangshan loess record shows a <100 Ma zircon age peak
316 (Fig. 6a). The zircon ages of the Mangshan loess are different from those of the middle reach of the Yellow
317 River (Fig. 6n), which are characterized by only one dominant Palaeozoic age peak at 200–350 Ma and two
318 old zircon populations at ca. 1850 Ma and ca. 2500 Ma (Stevens et al., 2013; Bird et al., 2015; Nie et al., 2015).
319 The Qin River and Yu River alluvial fan zircon ages (Figs. 6i and 6j), considered to be representative of the
320 provenance signal from the Taihang Mountains (Figs. 1a and 1b), are dominated by older zircon grains with
321 age peaks at ca. 1850 Ma and ca. 2500 Ma, respectively. The Yiluo River sample – representing the Qinling
322 Mountains source (Figs. 1a and 1b) – displays similar double-age peaks at 200–350 Ma and 350–550 Ma (Fig.
323 6d) as the Mangshan loess samples, with an additional younger age peak in the range of 60–180 Ma and a
324 distinct peak at ca. 2300 Ma, not present in the Mangshan loess. The Wei River, with its course north of the
325 Qinling Mountains (Fig. 1b), forms an important tributary of the Yellow River and carries sediments that are
326 dominated by the 200–300 Ma and especially the 350–550 Ma age populations (Fig. 6e). Notably, zircon grains
327 of early Cenozoic age (0–60 Ma) are also present in the Wei River sediments. Zircon ages of sediments from
328 the Sanmen Gorge (Figs. 1a and 1b), including Sanmen palaeolake sediments (Fig. 6h), fluvial sands of the
329 Sanmen Formation (Fig. 6f) and from the third and oldest river terrace (T3) of the Yellow River in Huangdigou

330 (HDG) of the Sanmen Gorge (Fig. 6g) (Wu et al., 1999) – all exhibit dominant peaks corresponding to 200–
331 350 Ma and 350–550 Ma, with two minor peaks at ca. 1850 Ma and ca. 2500 Ma.

332 Non-metric Multi-Dimensional Scaling (MDS) maps (Vermeesch, 2013; Vermeesch et al., 2016) are used
333 to visualize the (dis-) similarities between the zircon age profiles of Mangshan loess, CLP loess, Yellow River
334 sediments and other potential contributors to the Mangshan loess (Fig. 7). The Mangshan loess plots close to
335 the samples from the upper reach of the Yellow River and the CLP loess. The samples from the lower reach
336 of the Yellow River shows a close link with both the Wei River and the Yiluo River samples. The samples of
337 the Sanmen palaeolake and the river terraces in the Sanmen Gorge are located very close to each other. The
338 Qin River sample and the Yellow River middle reach sample lie close to each other but plot away from other
339 samples in the map.

340 **4. Discussion**

341 **4.1 Genetic interpretation of the Mangshan loess end-member model**

342 The end-member modelling results of the Mangshan loess-palaeosol grain-size record show that the
343 sediments are well described as a mixture of three different dust components which are comparable to the
344 average mixing model of the CLP (Figs. 3c and 3e) (Prins et al., 2007; Prins and Vriend, 2007). Prins et al.
345 (2007) interpreted the CLP loess-palaeosol records contain two different types of aeolian dust supplied from
346 two distinct source areas and/or reflecting different sediment transport-deposition process. The sandy (EM-1)
347 and silty (EM-2) loess components represent the coarse dust fraction transported by low level continental
348 northwesterly monsoonal winds via modified saltation and short term suspension processes over relative short
349 transport distances during major dust outbreaks. By contrast, the clayey loess component (EM-3) reflects a
350 fine dust component distributed over longer distances by long-term suspension processes. The fact that EM-1
351 and EM-2 are dominant in the Mangshan sequences indicates that most of the Mangshan loess was supplied
352 from a proximal source during major dust events. The independent mixing model of the last glacial-interglacial
353 sequence (S0-L1-S1) of the MS2006 section (Prins et al., 2009) shows that the clayey or ‘fine silty’ EM3 of
354 the upper Mangshan sequence (with the mode at 32 μm ; Fig. 3d) is coarser than EM3 of the composite MS2006
355 section (mode at 26 μm ; Fig. 3c). This observation suggests that the new mixing model allows better to make
356 the distinction between the two types of dust supply patterns also observed on the CLP.

357 According to the equation $MAR = 1.10F_{cf} + 6.12 (F_{cf} = F_{EM-1} + F_{EM-2})$ (section 3.3), the variation of F_{EM-3}
358 can be explained as a result of a “dilution effect” from the variation of the coarse flux F_{EM-1} and F_{EM-2} , with
359 high EM-3 content reflecting low MARs and low EM-3 content reflecting high MARs. Thus, variations in dust
360 flux during glacial and interglacials are expressed by relative high EM-3 content in palaeosol units (S1, S2,
361 S3, S4 and S5) and lower relative EM-3 content in loess units (L1, L2, L3, L4 and L5, Fig. 2c). The relative
362 content of fine EM-3 in loess units L1, L2 and L3 in the MS2006 section is significantly lower than in the
363 loess units L4, L5 and L6 in the lower part of the sequence, because of a higher input of the coarse dust fraction
364 of EM-1 and EM-2 ($F_{EM-1} + F_{EM-2}$) in the loess units L1, L2 and L3, particularly in L1 and L2. This is due to a
365 dramatically increased dust accumulation rate in the MS2006 section since the deposition of L3 (243–280 ka).
366 It is noticeable that both median grain size and proportions of modelled end members of the Mangshan record
367 fluctuate more frequently in the loess units L1 and L2 and palaeosol unit S1 (Fig.2). This observation might
368 reflect more variable climatic conditions during the last two glacial-interglacial cycles.

369 The results show a downwind (N-S) thinning and fining trend in grain-size (Fig. S5). In addition, the
370 proportion of modelled coarse fraction EM-1 also decreases from the MS2006 section to the more distal
371 MS2008W and MS2008E (Fig. S1). This spatial grain-size and EM-1 distribution pattern is also observed over
372 a much larger scale across the CLP (Prins et al., 2007; Prins and Vriend, 2007). Although this N-S downwind
373 pattern here is observed over a relatively small spatial scale (2.0-2.7 km), it likely suggests a proximal source
374 region to the north of the MLP, i.e the Yellow River floodplain for the Mangshan loess deposits, and indicates
375 that the near-surface northwesterly/northerly winter monsoon winds are responsible for the transportation of
376 the dust from the source area to the Mangshan Plateau.

377 4.2 Provenance signals of the Mangshan loess sequences

378 The overall zircon age distributions (Fig. 6) and the MDS map (Fig. 7) show that the Mangshan loess
379 deposits resemble those of the sediments from the upper and lower reaches of the Yellow River and of the CLP
380 loess. A genetic link between Yellow River sediments and CLP loess has already been proposed in previous
381 studies (Stevens et al., 2013; Bird et al., 2015; Nie et al., 2015). These studies suggested that the deposits on
382 the CLP are largely derived from the northeastern Tibetan Plateau (NTP) carried by the Yellow River and later
383 reworked by aeolian processes. A recent study by Licht et al. (2016) indicated further that the reworked Yellow

384 River sediments account for 60–70% of the supply to the CLP dust. The end member modelling on the grain-
385 size distribution of the Mangshan loess deposits imply that the coarse dust fractions EM1 and EM2 are derived
386 from the Yellow River floodplain just north of the MLP. Together with the zircon U-Pb age spectra this seems
387 to indicate that the Mangshan loess deposits have been constantly supplied by the lower Yellow River
388 floodplain since L9.

389 Sediments of the upper reach of the Yellow River exhibit two distinct Palaeozoic zircon age populations
390 at 200–350 Ma and 350–550 Ma matching the NTP provenance signatures. The lower reach of the Yellow
391 River, after the confluence of Wei River and Yiluo River, shows a similar zircon age distribution as the upper
392 reach of the Yellow River (Figs. 6m and 6o). According to Nie et al. (2015), this indicates the effects of the
393 Wei River and the Yiluo River (Fig.1b) which have brought abundant Phanerozoic zircon grains with a double
394 peak (200–550 Ma) from the Weihe Basin and Qinling Mountains to the lower Yellow River (Fig. 6), resulting
395 in a similar signal between the lower and upper reaches despite different source admixtures. In contrast, the
396 sediments of the middle reach of the Yellow River are distinctive from the upper and lower reaches of the river
397 by displaying a prominent single peak Palaeozoic age population at ca. 300 Ma (Fig. 6n) and two old
398 Proterozoic age populations at ca.1850 Ma and ca.2500 Ma. These ages are more similar to the source signals
399 of the Cretaceous bedrock and Northern China Craton, inherited from a river incision through Jihshan and
400 Sanmen canyons (Stevens et al., 2013; Bird et al., 2015; Nie et al., 2015; Zhang et al., 2016). The contribution
401 of debris from the Taihang Mountains to the lower Yellow River floodplain seems not significant as only a
402 subdued component of the older zircon age population (1800–3000 Ma) has been seen in the lower reach of
403 the Yellow River sediments.

404 Samples collected in the Sanmen Gorge area (Fig. 1) include fluvial sands of the Sanmen Formation (SM-
405 Fm) with an age of 1.4 Ma (Kong et al., 2014), lacustrine sands (SM-lake) from the upper Sanmen palaeolake,
406 and river terrace sands (SM-T3) from the third terrace in Huangdigou (HDG, Fig.1b) in the Sanmen Gorge .
407 (Jiang et al., 2007). Sediments of the Sanmen Formation (Fig. 6f) show similar zircon age distribution pattern
408 as that of the middle reach of the Yellow River, which has zircon age peaks at ca. 280 Ma, ca. 1850 Ma and
409 ca. 2500 Ma. This suggests that the Yellow River has flowed through Sanmen Gorge since 1.4 Ma (Kong et
410 al., 2014). The lacustrine sands of the Sanmen palaeolake (Fig. 6h) and the river terrace sample (SM-T3, Fig.6g)
411 also display a dominant zircon age peak at ca. 280 Ma, suggesting a major contribution of the middle reach of

the Yellow River sediments. However, it is interesting to note that compared to the other samples from the Sanmen Gorge, the zircon age of the Sanmen palaeolake (SM-lake, Fig. 6h) and the river terrace sample (SM-T3, Fig.6g) show an increase in the 350–550 Ma age population which is the dominant age component in sediments from the Wei River (Fig. 6e). Consistent with the age distribution patterns, the Sanmen palaeolake sample also lies close to the Wei River in the MDS map (Fig. 7). Based on the zircon provenance signal, Kong et al. (2014) concluded further that the Wei River primarily flowed through the Sanmen Gorge 1.3–1.5 Ma ago and then was followed by the Yellow River, which started to flow through the gorge from 1.3–1.4 Ma ago. As the time range provided by Kong et al. (2014) for the Yellow River and Wei River running through Sanmen Gorge largely overlap, it seems likely that both rivers flowed into the Sanmen palaeolake before it drained through Sanmen Gorge at around 1.3 Ma.

4.3 Implications from the sedimentological and provenance variations of the Mangshan loess-palaeosol sequences

The zircon U-Pb age data reveal that provenance signals of the upper and the lower reaches of the Yellow River dominate the Mangshan loess deposits from L9 to L1 (~900 ka to 15 ka), implying that the lower Yellow River floodplain has consistently served as the main source supply for the MLP since 900 ka. This conclusion is in agreement with results from Pan (2005), Hu et al. (2012), Kong et al. (2014) and Hu et al. (2017), showing that the Yellow River has flown through Sanmen Gorge at least since the early Pleistocene. Therefore, the observed dramatic increases in Mangshan Loess grain size and sedimentation rate since S2 could result from (1) approaching of the source area or extension of the floodplain of the lower Yellow River, (2) an increased sediment supply from the lower Yellow River, or (3) an increased wind strength. Because the sedimentation rate and grain size below and above palaeosol unit S2 in the loess-palaeosol sequences from the CLP (Vandenberghe et al., 1997; Ding et al., 2002; Sun, 2004; Sun and An, 2005; Sun et al., 2006) do not show a similar abrupt change as demonstrated in the Mangshan loess (Zheng et al., 2007), it is unlikely that winter monsoon intensity related wind strength changes played an important role. Thus, the sudden shift in sedimentology in the Mangshan sequence since 240 ka most probably arose from either the increased sediment supply of the lower Yellow River or the advance of its floodplain towards the MLP, or both.

Hu et al. (2012) found that the average incision rate of the Yellow River to the north of the Weihe Basin (Fig. 1b) increased dramatically since 240 ka as a result of local tectonic uplift. A recent study of the sedimentological infill of the Weihe Basin also suggests an increased incision rate of the fluvial system at approximately 240 ka resulting from enhanced tectonic activities (Rits et al., 2017). The finding suggests that differential tectonic movements in the Weihe Basin resulted in an increased sediment transfer propagating through the Sanmen Gorge. As a consequence, a wider alluvial fan system formed to the east of the Sanmen Gorge, next to the MLP (Fig. 1b). As this fan system served as a primary source for the MLP from ~240 ka onwards (Zheng et al., 2007), the accumulation rate on the plateau would have increased dramatically. An additional tectonic explanation for the observed sedimentation rate increase is the migration of the lower Yellow River floodplain towards the Mangshan Plateau. The high scarp present along the northern and eastern limits of the MLP is the result of lateral erosion by river action, and evidences a southward migration of the lower Yellow River floodplain during the late Pleistocene. The migration provided a more proximal dust source area for the Mangshan loess deposits. The southward migration are probably induced by subtle vertical motions related to NW-SE directed faults in the subsurface (Hao et al., 2008; Zhang et al., 2008).

5. Conclusions

The Mangshan loess-palaeosol record provides a high-resolution archive of dust supply from the Yellow River floodplain. The high accumulation rates, the coarse-grained character of the loess, and the distinct thinning and fining of the loess deposits in a north-south direction, clearly indicate increased coarser dust supply from the Yellow River floodplain to the Mangshan Plateau during the last two glacial-interglacial intervals. An independent provenance record by single grain zircon U-Pb ages confirms the lower Yellow River floodplain as the likely main source for the Mangshan dust deposits, at least since loess unit L9 (900 ka). This implies that the Yellow River incised the Sanmen Gorge before 900 ka (~MIS 22, i.e. late Early Pleistocene). The dramatic sedimentological change in the Mangshan sequences above S2 most likely originates from the tectonic activity in the Weihe Basin since 240 ka. This resulted in accelerated incision rate of the fluvial systems and associated release of eroded material through Sanmen Gorge towards the lower reach of the Yellow River floodplain. Meanwhile, a simultaneous southward migration and lateral erosion of

464 the (lower) Yellow River resulted in a more proximal location of the dust source area, all contributing to
465 increased loess deposition on the Mangshan Plateau during the late Pleistocene.

466

467 **Acknowledgements**

468 We thank Patrick Bacon, Xiangtong Huang, Ilse Kamerling, Simon Troelstra, Noortje Vromans, Ke Wang,
469 Wieske Wentink, Hugo Wester, Wouter Wester, Bin Zhou and local farmers from the Liugou Village for their
470 help with sampling the Mangshan loess-palaeosol sequences in 2006 and 2008. Roel van Elsas is thanked for
471 supervision of heavy mineral separation. MAP and CJB thank the Royal Netherlands Academy of Arts and
472 Sciences (KNAW) for financial support of these field projects. YS thanks the Doctoral Program of Geosciences
473 of University of Helsinki for travel funding. BW acknowledges the Natural Science Foundation of China
474 (Project Number 41602182) for financial support of the field work. We thank Xiaoping Yang for editorial
475 work and two anonymous reviewers for their helpful and constructive reviews of the manuscript. This research
476 is supported by the Academy of Finland grant to AK (Projects Numbers 257850, 264935 and 292827) .

477 **References**

478

479 AFSOM, 1988. Active Fault System Around Ordos Massif. Seismological Press, Beijing.

480 An, Z.S., Kutzbach, J.E., Prell, W.L., Porter, S.C., 2001. Evolution of Asian monsoons and phased uplift of
481 the Himalaya-Tibetan plateau since Late Miocene times. *Nature* 411, 62-66.

482 Berger, A.L., 1978. Long-Term Variations of Caloric Insolation Resulting from the Earth's Orbital Elements.
483 *Quat. Res.* 9, 139-167.

484 Bird, A., Stevens, T., Rittner, M., Vermeesch, P., Carter, A., Andò, S., Garzanti, E., Lu, H., Nie, J., Zeng, L.,
485 Zhang, H., Xu, Z., 2015. Quaternary dust source variation across the Chinese Loess Plateau. *Palaeogeogr.*
486 *Palaeoclimatol. Palaeoecol.* 435, 254-264.

487 Cheng, H., Edwards, R.L., Broecker, W.S., Denton, G.H., Kong, X., Wang, Y., Zhang, R., Wang, X., 2009.
488 Ice age terminations. *Science* 326, 248-252.

489 Cheng, H., Edwards, R.L., Sinha, A., Spotl, C., Yi, L., Chen, S., Kelly, M., Kathayat, G., Wang, X., Li, X.,
490 Kong, X., Wang, Y., Ning, Y., Zhang, H., 2016. The Asian monsoon over the past 640,000 years and ice age
491 terminations. *Nature* 534, 640-646.

492 Deplazes, G., Lückge, A., Stuut, J.-B.W., Pätzold, J., Kuhlmann, H., Husson, D., Fant, M., Haug, G.H., 2014.
 493 Weakening and strengthening of the Indian monsoon during Heinrich events and Dansgaard-Oeschger
 494 oscillations. *Paleoceanography* 29, 99-114.

495 Derbyshire, E., Meng, X., Kemp, R.A., 1998. Provenance, transport and characteristics of modern aeolian dust
 496 in western Gansu Province, China, and interpretation of the Quaternary loess record. *J. Arid Environ.* 39, 497-
 497 516.

498 Ding, Z.L., Derbyshire, E., Yang, S.L., Sun, J.M., Liu, T.S., 2005. Stepwise expansion of desert environment
 499 across northern China in the past 3.5 Ma and implications for monsoon evolution. *Earth Planet. Sci. Lett.* 237,
 500 45-55.

501 Ding, Z.L., Derbyshire, E., Yang, S.L., Yu, Z.W., Xiong, S.F., Liu, T.S., 2002. Stacked 2.6-Ma grain size
 502 record from the Chinese loess based on five sections and correlation with the deep-sea $\delta^{18}\text{O}$ record.
 503 *Paleoceanography* 17, doi:10.1029/2001PA000725.

504 Hao, K., Tian, Q., Liu, B., Yin, G., 2008. Exploration of the Laoyachen fault, Zhengzhou of China, and its
 505 activity investigation. *Acta Seismol. Sin.* 30, 416-423.

506 Hu, Z., Pan, B., Bridgland, D., Vandenberghe, J., Guo, L., Fan, Y., Westaway, R., 2017. The linking of the
 507 upper-middle and lower reaches of the Yellow River as a result of fluvial entrenchment. *Quat. Sci. Rev.* 166,
 508 324–338.

509 Hu, Z., Pan, B., Wang, J., Cao, B., Gao, H., 2012. Fluvial terrace formation in the eastern Fenwei Basin, China,
 510 during the past 1.2Ma as a combined archive of tectonics and climate change. *J Asian Earth Sci.* 60, 235-245.

511 Huang, C.C., Pang, J., Su, H., Li, S., Ge, B., 2009. Holocene environmental change inferred from the loess–
 512 palaeosol sequences adjacent to the floodplain of the Yellow River, China. *Quat. Sci. Rev.* 28, 2633-2646.

513 Ji, J.L., Zheng, H.B., Liu, R., Huang, X., Jiang, F.C., 2004. Restudy on the stratigraphy of Mangshan loess.
 514 *Mar. Geol. Quat. Geol.* 24, 101-108.

515 Jiang, F., Fu, J., Wang, S., Sun, D., Zhao, Z., 2007. Formation of the Yellow River, inferred from loess–
 516 palaeosol sequence in Mangshan and lacustrine sediments in Sanmen Gorge, China. *Quat. Int.* 175, 62-70.

517 Jiang, F.C., Wang, S.B., Zhao, Z.Z., Fu, J.L., 2004. Mangshan loess in Central China and the paleomonsoon
 518 variations since the last interglaciation. *Acta Geol. Sin.* 78, 813-819, doi:10.1111/j.1755-6724.2004.tb00200.x.

519 Jiang, F.C., Wu, X.H., Sun, D.H., Wang, S.M., An, Z.S., Tian, G.Q., Liu, K., Yin, W.D., Xue, B., 1998. On
 520 Mangshan loess stratigraphy in China central plains. *J. Geomech.* 4, 90-97.

521 Kohfeld, K.E., Harrison, S.P., 2003. Glacial-interglacial changes in dust deposition on the Chinese Loess
522 Plateau. *Quat. Sci. Rev.* 22, 1859-1878.

523 Konert, M., Vandenberghe, J., 1997. Comparison of laser grain size analysis with pipette and sieve analysis: a
524 solution for the underestimation of the clay fraction. *Sedimentology* 44, 523-535.

525 Kong, P., Jia, J., Zheng, Y., 2014. Time constraints for the Yellow River traversing the Sanmen Gorge.
526 *Geochem. Geophys. Geosyst* 15, 395-407, doi:10.1002/2013GC004912.

527 Kukla, G., 1987. Loess stratigraphy in central China. *Quat. Sci. Rev.* 6, 191-219.

528 Li, B., Sun, D., Xu, W., Wang, F., Liang, B., Ma, Z., Wang, X., Li, Z., Chen, F., 2017. Paleomagnetic
529 chronology and paleoenvironmental records from drill cores from the Hetao Basin and their implications for
530 the formation of the Hobq Desert and the Yellow River. *Quat. Sci. Rev.* 156, 69-89.

531 Licht, A., Pullen, A., Kapp, P., Abell, J., Giesler, N., 2016. Eolian cannibalism: Reworked loess and fluvial
532 sediment as the main sources of the Chinese Loess Plateau. *Geol. Soc. Am. Bull.* B31375.31371, doi:
533 10.1130/B31375.1.

534 Lin, A., Yang, Z., Sun, Z., Yang, T., 2001. How and when did the Yellow River develop its square bend?
535 *Geology* 29, 951-954.

536 Lisiecki, L.E., Raymo, M.E., 2005. A Pliocene-Pleistocene stack of 57 globally distributed benthic $\delta^{18}\text{O}$
537 records. *Paleoceanography* 20, doi:10.1029/2004PA001071.

538 Liu, T., Ding, Z., 1998. Chinese Loess and the Paleomonsoon. *Annu. Rev. Earth Planet. Sci.* 26, 111-145.

539 Liu, T.S., 1985. *Loess and the Environment*. China Ocean Press, Beijing.

540 Lu, H.Y., Sun, D.H., 2000. Pathways of dust input to the Chinese Loess Plateau during the last glacial and
541 interglacial periods. *Catena* 40, 251-261.

542 Nie, J., Stevens, T., Rittner, M., Stockli, D., Garzanti, E., Limonta, M., Bird, A., Ando, S., Vermeesch, P.,
543 Saylor, J., Lu, H., Breecker, D., Hu, X., Liu, S., Resentini, A., Vezzoli, G., Peng, W., Carter, A., Ji, S., Pan,
544 B., 2015. Loess Plateau storage of Northeastern Tibetan Plateau-derived Yellow River sediment. *Nature*
545 *Commun.* 6, 8511, doi:10.1038/ncomms9511.

546 Nugteren, G., Vandenberghe, J., 2004. Spatial climatic variability on the Central Loess Plateau (China) as
547 recorded by grain size for the last 250 kyr. *Global Planet. Change* 41, 185-206.

548 Pan, B., 2005. Paleomagnetic dating of the topmost terrace in Kouma, Henan and its indication to the Yellow
549 River's running through Sanmen Gorges. *Chinese Sci. Bull.* 50, 657-664.

550 Pan, B., Hu, Z., Wang, J., Vandenberghe, J., Hu, X., 2011. A magnetostratigraphic record of landscape
551 development in the eastern Ordos Plateau, China: Transition from Late Miocene and Early Pliocene stacked
552 sedimentation to Late Pliocene and Quaternary uplift and incision by the Yellow River. *Geomorphology* 125,
553 225-238.

554 Peterse, F., Prins, M.A., Beets, C.J., Troelstra, S.R., Zheng, H., Gu, Z., Schouten, S., Damsté, J.S.S., 2011.
555 Decoupled warming and monsoon precipitation in East Asia over the last deglaciation. *Earth Planet. Sci. Lett.*
556 301, 256-264.

557 Porter, S.C., 2001. Chinese loess record of monsoon climate during the last glacial–interglacial cycle. *Earth-*
558 *Sci. Rev.* 54, 115-128.

559 Prins, M.A., Postma, G., Weltje, G.J., 2000. Controls on terrigenous sediment supply to the Arabian Sea during
560 the late Quaternary: the Makran continental slope. *Mar. Geol.* 169, 351-371.

561 Prins, M.A., Vriend, M., 2007. Glacial and interglacial eolian dust dispersal patterns across the Chinese Loess
562 Plateau inferred from decomposed loess grain-size records. *Geochem. Geophys. Geosyst.* 8, Q07Q05,
563 doi:10.1029/2006GC001563.

564 Prins, M.A., Vriend, M., Nugteren, G., Vandenberghe, J., Lu, H., Zheng, H., Jan Weltje, G., 2007. Late
565 Quaternary aeolian dust input variability on the Chinese Loess Plateau: inferences from unmixing of loess
566 grain-size records. *Quat. Sci. Rev.* 26, 230-242.

567 Prins, M.A., Weltje, G.J., 1999. End-member modeling of siliciclastic grain-size distributions: the Late
568 Quaternary record of eolian and fluvial sediment supply to the Arabian Sea and its paleoclimatic significance.
569 In: Harbaugh, J., Watney, L., Rankey, G., Slingerland, R., Goldstein, R., Franseen, E. (Eds.), *Numerical*
570 *Experiments in Stratigraphy: Recent Advances in Stratigraphic and Sedimentologic Computer Simulations.*
571 *SEPM (Society for Sedimentary Geology) Special Publication* 62, pp. 91-111.

572 Prins, M.A., Zheng, H., Beets, K., Troelstra, S., Bacon, P., Kamerling, I., Wester, W., Konert, M., Huang, X.,
573 Ke, W., Vandenberghe, J., 2009. Dust supply from river floodplains: the case of the lower Huang He (Yellow
574 River) recorded in a loess-palaeosol sequence from the Mangshan Plateau. *J. Quat. Sci.* 24, 75-84.

575 Pye, K., 1995. The nature, origin and accumulation of loess. *Quat. Sci. Rev.* 14, 653-667.

576 Qiu, F., Zhou, L., 2015. A new luminescence chronology for the Mangshan loess-palaeosol sequence on the
577 southern bank of the Yellow River in Henan, central China. *Quat. Geochronol.* 30, 24-33.

578 Rits, D.S., van Balen, R.T., Prins, M.A., Zheng, H., 2017. Evolution of the alluvial fans of the Luo River in
579 the Weihe Basin, central China, controlled by faulting and climate change - A reevaluation of the
580 paleogeographical setting of Dali Man site. *Quat. Sci. Rev.* 166, 339-351.

581 Shang, Y., Beets, C.J., Tang, H., Prins, M.A., Lahaye, Y., van Elsas, R., Sukselainen, L., Kaakinen, A., 2016.
582 Variations in the provenance of the late Neogene Red Clay deposits in northern China. *Earth Planet. Sci. Lett.*
583 439, 88-100.

584 Shang, Y., Kaakinen, A., Beets, C.J., Prins, M.A., in press. Aeolian silt transport processes as fingerprinted by
585 dynamic image analysis of the grain size and shape characteristics of Chinese loess and Red Clay deposits.
586 *Sediment. Geol.* doi:10.1016/j.sedgeo.2017.12.001.

587 Stacey, J.S., Kramers, J.D. 1975. Approximation of terrestrial lead isotope evolution by a two-stage model.
588 *Earth Planet. Sci. Lett.* 26, 207-221.

589 Stevens, T., Carter, A., Watson, T.P., Vermeesch, P., Andò, S., Bird, A.F., Lu, H., Garzanti, E., Cottam, M.A.,
590 Sevastjanova, I., 2013. Genetic linkage between the Yellow River, the Mu Us desert and the Chinese Loess
591 Plateau. *Quat. Sci. Rev.* 78, 355-368.

592 Stuut, J.-B.W., Prins, M.A., Schneider, R.R., Weltje, G.J., Jansen, J.H.F., Postma, G., 2002. A 300-kyr record
593 of aridity and wind strength in southwestern Africa: inferences from grain-size distributions of sediments on
594 Walvis Ridge, SE Atlantic. *Mar. Geol.* 180, 221-233.

595 Stuut, J.-B.W., Temmesfeld, F., De Deckker, P., 2014. A 550 ka record of aeolian activity near North West
596 Cape, Australia: inferences from grain-size distributions and bulk chemistry of SE Indian Ocean deep-sea
597 sediments. *Quat. Sci. Rev.* 83, 83-94.

598 Sun, D.H., 2004. Monsoon and westerly circulation changes recorded in the late Cenozoic aeolian sequences
599 of Northern China. *Global Planet. Change* 41, 63-80.

600 Sun, D.H., Bloemendal, J., Rea, D.K., An, Z., Vandenberghe, J., Lu, H., Su, R., Liu, T., 2004. Bimodal grain-
601 size distribution of Chinese loess, and its palaeoclimatic implications. *Catena* 55, 325-340.

602 Sun, D.H., Bloemendal, J., Rea, D.K., Vandenberghe, J., Jiang, F., An, Z., Su, R., 2002. Grain-size distribution
603 function of polymodal sediments in hydraulic and aeolian environments, and numerical partitioning of the
604 sedimentary components. *Sediment. Geol.* 152, 263-277.

605 Sun, J., 2002. Provenance of loess material and formation of loess deposits on the Chinese Loess Plateau. *Earth*
606 *Planet. Sci. Lett.* 203, 845-859.

607 Sun, Y., An, Z., 2005. Late Pliocene-Pleistocene changes in mass accumulation rates of eolian deposits on the
608 central Chinese Loess Plateau. *J. Geophys. Res. Atmos.* 110, D23101.

609 Sun, Y., Chen, J., Clemens, S.C., Liu, Q., Ji, J., Tada, R., 2006. East Asian monsoon variability over the last
610 seven glacial cycles recorded by a loess sequence from the northwestern Chinese Loess Plateau. *Geochem.*
611 *Geophys. Geosyst.* 7, Q12Q02, doi:10.1029/2006GC001287.

612 Van Achterbergh, E., Ryan C., Jackson, S., Griffin, W., 2001. Data reduction software for LA-ICP-MS, in: P.,
613 S. (Ed.), *Laser-Ablation ICPMS in the Earth Sciences – Principles and applications*. Mineralogical Association
614 of Canada short course series, St John, Newfoundland, pp. 239-243.

615 Vandenberghe, J., An, Z.S., Nugteren, G., Lu, H.Y., Van Huissteden, K., 1997. New absolute time scale for
616 the Quaternary climate in the Chinese loess region by grain-size analysis. *Geology* 25, 35-38.

617 Vermeesch, P., 2013. Multi-sample comparison of detrital age distributions. *Chem. Geol.* 341, 140-146.

618 Vermeesch, P., Resentini, A., Garzanti, E., 2016. An R package for statistical provenance analysis. *Sediment.*
619 *Geol.* 336, 14-25.

620 Vriend, M., Prins, M.A., Buylaert, J.P., Vandenberghe, J., Lu, H.Y., 2011. Contrasting dust supply patterns
621 across the north-western Chinese Loess Plateau during the last glacial-interglacial cycle. *Quat. Int.* 240, 167-
622 180.

623 Wang, S., 2002. Sedimentary records of environmental evolution in the Sanmen Lake Basin and the Yellow
624 River running through the Sanmenxia Gorge eastward into the sea. *Sci. China, Ser. D Earth Sci.*, 45, 595-608.

625 Wang, S., Fu, B., Piao, S., Lü, Y., Ciais, P., Feng, X., Wang, Y., 2015. Reduced sediment transport in the
626 Yellow River due to anthropogenic changes. *Nature Geosci.* 9, 38-41.

627 Wang, S.B., Jiang, F.C., Fu, J.L., Li, C.Z., Cai, Y., Yao, H.T., Qiao, Y.S., Zhang, Z.S., Li, Y.J., 2013. Some
628 knowledge of the formation of the Yellow River. *Quat. Sci.* 33, 705-714.

629 Wang, Y., Cheng, H., Edwards, R.L., Kong, X., Shao, X., Chen, S., Wu, J., Jiang, X., Wang, X., An, Z., 2008.
630 Millennial- and orbital-scale changes in the East Asian monsoon over the past 224,000 years. *Nature* 451,
631 1090-1093.

632 Weltje, G.J., 1997. End-member modeling of compositional data: Numerical-statistical algorithms for solving
633 the explicit mixing problem. *Math. Geol.* 29, 503-549.

634 Weltje, G.J., Prins, M.A., 2003. Muddled or mixed? Inferring palaeoclimate from size distributions of deep-
635 sea clastics. *Sediment. Geol.* 162, 39-62.

636 Weltje, G.J., Prins, M.A., 2007. Genetically meaningful decomposition of grain-size distributions. *Sediment.*
637 *Geol.* 202, 409-424.

638 Wu, X., Jiang, F., Xiao, H., Xue, B., Sun, D., 1999. Mangshan loess on China' s Central Plain and its response
639 to tectonic movement and climate*. *Sci. China, Ser. D Earth Sci.*, 42, 465-473, doi:10.1007/BF02875240.

640 Xiao, J., Chang, Z., Si, B., Qin, X., Itoh, S., Lomtatidze, Z., 2009. Partitioning of the grain-size components
641 of Dali Lake core sediments: evidence for lake-level changes during the Holocene. *J. Paleolimnol.* 42, 249-
642 260.

643 Xiao, J., Fan, J., Zhou, L., Zhai, D., Wen, R., Qin, X., 2013. A model for linking grain-size component to lake
644 level status of a modern clastic lake. *J. Asian Earth Sci.* 69, 149-158.

645 Zhang, H., Lu, H., Xu, X., Liu, X., Yang, T., Stevens, T., Bird, A., Xu, Z., Zhang, T., Lei, F., Feng, H., 2016.
646 Quantitative estimation of the contribution of dust sources to Chinese loess using detrital zircon U-Pb age
647 patterns. *J. Geophys. Res. Earth Surf.* 121, 2085-2099.

648 Zhang, J., Shen, J., Tian, Q., Shen, X., Zhang, X., Li, B., 2008. Relationship Between Mangshan Geomorphic
649 Scarp and Laoyachen Fault Activity. *Earthquake* 28, 121-127.

650 Zheng, H., Huang, X., Ji, J., Liu, R., Zeng, Q., Jiang, F., 2007. Ultra-high rates of loess sedimentation at
651 Zhengzhou since Stage 7: Implication for the Yellow River erosion of the Sanmen Gorge. *Geomorphology* 85,
652 131-142.

653

654 **Figures**

655 **Figure 1.** (a) Digital elevation model (DEM) map of northern China showing the Yellow River, Chinese Loess
656 Plateau (CLP) and Mangshan Loess Plateau (MLP). The grey arrows indicate the direction of winter monsoon
657 winds. The red letters indicate the boundary of upper, middle and lower reaches of the Yellow River. The
658 white triangles indicate the loess sections on the CLP referred to in the text and the white dots indicate the
659 previous published Yellow River samples of zircon U-Pb ages (Nie et al., 2015). Samples 1-11, 12-13 and 14-
660 15 are representative for the upper, middle and lower reaches of the Yellow River respectively. (b) Map
661 showing the zircon sampling sites in the Weihe Basin, middle and lower reaches of the Yellow River. The red
662 triangle marks the MLP. The black dots show samples collected within this study while the white dots represent
663 samples cited in Kong et al. (2014). Detailed sample description is presented in Table S3. HDG is abbreviation
664 for Huangdigou, where SM-T3 and SM-lake being collected. (c) Loess section MS2006, MS2008W and
665 MS2008E on the MLP of this study marked with red triangles. Figure (c) is produced with Google Earth.
666 Imagery © 2017 TerraMetrics. Data © Europa Technologies Ltd.

667
668 **Figure 2.** (a) Palaeosol (S0, S1 ... S5) and loess (L1, L2 ... L6) stratigraphy in section MS2006 with black
669 dots indicating the levels of zircon samples; (b) median grain size distribution of the MS2006 section, (c) the
670 proportional contribution of the end members against depth. The age model of MS2006 is based on tuning to
671 the following ‘target curves’ (time series): (d) the oxygen-isotope composite record from Dongge, Sanbao and
672 Hulu caves in central China (Cheng et al., 2009; Cheng et al., 2016; Wang et al., 2008), here superimposed on
673 the summer (21 July) insolation at 65°N (Berger, 1978), and (e) the stacked marine benthic oxygen-isotope
674 record (Lisiecki and Raymo, 2005). The grey bars indicate palaeosol layers and corresponding interglacial
675 stages / marine isotope stages (MIS). The tie points linking the loess-palaeosol record to the target isotope
676 curves are listed in Table S1. Section MS2006 (a) is ~130 m thick and is a composite of record X (0-59 m;
677 Prins et al., 2009) and record Z’ (59-130 m; Zheng et al., 2007, see also Prins et al., 2009).

678
679 **Figure 3.** End-member modelling results of the composite MS2006 section. (a) The mean/median r^2 across
680 the full-size range as a function of the number of end members (q). (b) Coefficient of determination (r^2)
681 statistics for each size class for end-member models (EMM) with 2–5 end members. (c) Modelled end members

682 according to a three-end-member model representing sandy loess (EM-1, modal size $\sim 63 \mu\text{m}$), silty loess (EM-
683 2, modal size $\sim 37 \mu\text{m}$) and clayey loess (EM-3, modal size $\sim 26 \mu\text{m}$) for the composite MS2006 section in this
684 study, (d) the last glacial-interglacial sequence (S0-L1-S1) from the Mangshan Plateau (Prins et al., 2009) and
685 (e) the Chinese Loess Plateau (Prins and Vriend, 2007).

686

687 **Figure 4.** (a) Age-depth plot (dashed line) of the loess-palaeosol boundaries in the MS2006 section, based on
688 the correlation of the loess-palaeosol record to the target isotope curves shown in Fig. 2, and corresponding
689 linear sedimentation rate (LSR) estimates per loess and palaeosol unit (solid line). (b) Scatter plot of the ‘coarse
690 fraction’ flux (F_{cf} , i.e. flux of EM1 and EM2) *versus* the total mass-accumulation rate (MAR) of all major loess
691 (L1, L2 ... L5) and palaeosol units (S1, S2 ... S5) in MS2006 (and MS2008W, see legend). Linear regression
692 formula for the MS2006 dataset is shown. Data are listed in Table S1 and Table S2.

693

694 **Figure 5.** (a) Palaeosol (S0, S1 ... S5) and loess (L1, L2 ... L6) stratigraphy in section MS2006 with red dots
695 showing the level of samples for zircon U-Pb age analysis. (b) Zircon U-Pb age distributions of Mangshan
696 loess samples. The black lines are normalised probability density function plots (PDP); the orange shades are
697 Kernel Density Estimation (KDE) plots for different units; the open rectangles are age histograms. The blue
698 rectangle indicates the 0–100 Ma age population while grey rectangles mark the dominant age population 250–
699 350 and 350–550 Ma in the spectra.

700

701 **Figure 6.** Detrital zircon U-Pb ages for samples analysed within this study and the previously published dataset
702 for CLP loess (Bird et al., 2015), the Wei River (Kong et al., 2014), the Sanmen Formation (Kong et al., 2014)
703 and the Yellow River upper, middle and lower reaches (Nie et al., 2015). The samples’ locations are indicated
704 in Fig. 1 and detailed description are presented in Supplementary Table S3. MS-upper (6a) and MS-lower (6b)
705 are the combined zircon age dataset of samples from the upper part (L1-1, L1-2, L2-1 and L2-2) and lower
706 part (L5, L6 and L9) of the MS2006 section respectively. Note we excluded sample L3 from the combination
707 dataset because loess unit L3 is a transition period in the sedimentology of the stratigraphy. The black lines
708 are normalised probability density function plots (PDP); the colour shades are Kernel Density Estimation

709 (KDE) plots for different units and the open rectangles are age histograms. Vertical dash lines indicate the
710 major age peaks of the spectra.

711 **Figure 7.** Non-metric multi-dimensional scaling (MDS) map visualising the comparison between the
712 Mangshan zircon age dataset (MS-U and MS-L), the loess of the Chinese Loess Plateau (CLP Loess), and
713 zircon datasets of fluvial deposits of the Yellow River upper reach (YR-U), middle reach (YR-M) and lower
714 reach (YR-Ln, combination dataset of samples YR-L, YR-20 and YR-33 in Fig. 6), the Yiluo River (YL-R),
715 the Wei River (WR), the Qin River (QR) and samples of the Sanmen palaeolake (SM-lake) and fluvial sands
716 of the Sanmen Formation (SM-Fm). The stress value is 0.35%, indicating an “excellent fit” of the data. The
717 solid lines link the closest neighbours and the dashed lines the second closest neighbours.




Figure 1






Figure 2


(a)




(b)



(c) Composite MS2006 section



(d) MS2006 (S0-L1-S1)



(e) Chinese Loess Plateau




Figure 3





Figure 4




Figure 5




Figure 6




Figure 7

Supplementary material

Table S1. Depth and estimated ages of the stratigraphic boundaries between loess and palaeosol units in sections MS2006 and MS2008W

Stratigraphic boundary	MS2006 Depth (m)	MS2008W Depth (m)	Age (ka)
S0-L1	1.28	2.45	11.5 ^a
L1-S1	40.18	26.15	77.1 ^a
S1-L2	57.73	33.75	129 ^a
L2-S2	85.24		191.8 ^a
S2-L3	93.94		242.5 ^a
L3-S3	104.00		281 ^a
S3-L4	107.40		336 ^a
L4-S4	111.20		396 ^a
S4-L5	114.10		424 ^a
L5-S5	121.50		473 ^a
S5-L6	125.10		624 ^a

^aAges according to speleothem oxygen isotope records of Cheng et al. (2016) and references therein.

Table S2. Data of sections MS2006 and MS2008W used to calculate the end-member specific fluxes per loess/palaeosol unit, including sedimentation rate (SR), mass-accumulation rate (MAR) and end-member proportions (p_{EM-x})

Section	Unit	Thickness (m)	SR (cm/ka)	MAR (g/cm ² /ka)	p_{EM-1}	p_{EM-2}	p_{EM-3}
MS2006	S0	1.28	11	16	0.18	0.41	0.40
	L1	38.90	59	88	0.45	0.41	0.14
	S1	17.55	34	50	0.17	0.50	0.33
	L2	27.52	44	65	0.37	0.46	0.17
	S2	8.70	17	25	0.04	0.46	0.50
	L3	10.06	26	39	0.09	0.77	0.14
	S3	3.40	6	9	0.04	0.34	0.62
	L4	3.80	6	9	0.01	0.62	0.37
	S4	2.90	10	15	0.06	0.37	0.57
	L5	7.40	15	22	0.12	0.51	0.37
	S5	3.60	2	4	0.03	0.29	0.68
MS2008 W	S0	2.45	21	32	0.07	0.55	0.38
	L1	23.70	36	53	0.20	0.62	0.18
	S1	7.60	15	22	0.04	0.55	0.41

Table S3. Description of location and nature of the zircons U-Pb samples used in this study

Sample name	Coordinates		Nature	Reference
	N (°)	E(°)		
MS-L1-1	34.96°	113.37°	Loess unit L1, MS2006 section	This study
MS-L1-2	34.96°	113.37°	Loess uit L1, MS2006 section	This study
MS-L2-1	34.96°	113.37°	Loess unit L2, MS2006 section	This study
MS-L2-2	34.96°	113.37°	Loess unit L2, resampled from Mangshan	This study
MS-L3	34.96°	113.37°	Loess unit L3, MS2006 section	This study
MS-L5	34.96°	113.37°	Loess unit L5, MS2006 section	This study
MS-L6	34.96°	113.37°	Loess unit L6, MS2006 section	This study
MS-L9	34.97°	113.37°	Loess unit L9, resampled from Mangshan	This study
Qin River	35.14°	112.79°	River sediment	This study
Yiluo River	34.71°	112.59°	River sediment	This study
YR-20	34.85°	112.62°	Yellow River sediment, near Mengjin county	This study
YR-33	34.98°	113.38°	Yellow River sediment, near Mangshan	This study
SM-Lake	34.87°	111.3°	Sanmen palaeolake, fluvial sand	This study
SM-T3	34.82°	111.28°	River terrace sediment in Huangdigou	This study
Yu River	35.43°	113.45°	Modern alluvial fan sand	This study
Wei River	34.45°	109.52°	River sediment, Wei River	Samplpe SM18 in Kong et al., 2014
SM Fm	34.7°	110.29°	Fluvial sands from Sanmen Formation	Sample SM12 and 13 in Kong et al., 2014
CLP loess			Loess samples from Beiguoyuan and Lingtai	Bird et al., 2015
YR-U			Yellow River sediment, upper reach	Nie et al., 2015
YR-M			Yellow River sediment, middle reach	Nie et al., 2015
YR-L			Yellow River sediment, lower reach	Nie et al., 2015

Figures




Figure S1. Schematic N-S transect from the Yellow River floodplain to sections MS2006 and MS2008W on the Manghan Loess Plateau. Large dots with error bars indicate altitude measurements (from GPS) at the three sites (top of sections). The present-day geomorphology of the plateau is characterized by a natural N-S sloping southern flank with an angle of $\sim 1.15^\circ$ (gradient of 40 m over 2 km horizontal distance). Solid correlation lines between the two loess sections highlight the N-S thinning of loess unit L1 (38%) and palaeosol unit S1 (55%). Extrapolation of these ‘thinning factors’ to the successive L2-S2 sequence in MS2008W (dashed correlation lines) suggests that the wedge-shape geometry of the loess-palaeosol sediment package exists since the formation of palaeosol S2. B/M is the Bruhnes/Matuyama boundary which is found in the lower part of L8 in the MS2006 section (Zheng et al., 2007).




Figure S2. Carbonate, organic matter and median grain size compared to palaeosol (S0, S1 ... S5) and loess (L1, L2 ... L6) units in section MS2006 (composite of records X and Z'). Note the overall change in character of the loess-palaeosol layers from base to top: the palaeosol and loess layers in the lower part of the profile (S3-S5, L3-L6) are relative thin, fine-grained and organic-rich, whereas the palaeosol (S0-S2) and loess layers (L1 and L2) in the upper part of the sequence are significantly thicker, coarser-grained and contain less organic matter. The grey rectangle indicates the overlap part of record X and Z' (Fig. S3).




Figure S3. Median grain size, organic matter and carbonate content in the interval 36-60 m of section MS2006, i.e. the interval where records X (solid lines with symbols) and Z' (solid lines without symbols) overlap. The stratigraphic profiles in records X and Z' are very similar, despite the fact that the records have independent depth scales (see Prins et al., 2009 for details).




Figure S4. Magnetic susceptibility, carbonate content and median grain size, compared to palaeosol (S0, S1) and loess (L1, L2) units in (the upper part of) section MS2006 (record X), MS2008W and MS2008E. Dashed lines correlate lithological (and marine-isotope stage) boundaries. Note that section MS2006 and the two MS2008 sections are separated by ~3 km. Profiles MS2006 and MS2008W are lined up with respect to the top of their S0 units (ground level) to highlight the N-S thinning of the S0-L1-S1 sequence; profile MS2008E is lined up with respect to the S1 unit in profile MS2008W, showing a local (palaeo-) relief of ~3.7 meter. Figure modified after Peterse et al. (2011).




Figure S5. Proportional contribution of the end members and magnetic susceptibility compared to palaeosol (S0, S1) and loess (L1, L2) stratigraphy in (the upper part of) section MS2006 (record X), MS2008W and MS2008E. Dashed correlation lines coincide with lithological (and marine-isotope stage) boundaries. Note the significant lateral changes between sections MS2006 and MS2008W in L1 and S1 layer thicknesses and corresponding shifts in the end-member mixing coefficients.

References:

- Bird, A., Stevens, T., Rittner, M., Vermeesch, P., Carter, A., Andò, S., Garzanti, E., Lu, H., Nie, J., Zeng, L., Zhang, H., Xu, Z., 2015. Quaternary dust source variation across the Chinese Loess Plateau. *Palaeogeogr. Palaeoclimatol. Palaeoecol.* 435, 254-264.
- Cheng, H., Edwards, R.L., Sinha, A., Spotl, C., Yi, L., Chen, S., Kelly, M., Kathayat, G., Wang, X., Li, X., Kong, X., Wang, Y., Ning, Y., Zhang, H., 2016. The Asian monsoon over the past 640,000 years and ice age terminations. *Nature* 534, 640-646.
- Kong, P., Jia, J., Zheng, Y., 2014. Time constraints for the Yellow River traversing the Sanmen Gorge. *Geochemistry, Geophysics, Geosystems* 15, 395-407.
- Nie, J., Stevens, T., Rittner, M., Stockli, D., Garzanti, E., Limonta, M., Bird, A., Ando, S., Vermeesch, P., Saylor, J., Lu, H., Breecker, D., Hu, X., Liu, S., Resentini, A., Vezzoli, G., Peng, W., Carter, A., Ji, S., Pan, B., 2015. Loess Plateau storage of Northeastern Tibetan Plateau-derived Yellow River sediment. *Nature commun.* 6, 8511, doi:10.1038/ncomms9511.

Peterse, F., Prins, M.A., Beets, C.J., Troelstra, S.R., Zheng, H., Gu, Z., Schouten, S., Damsté, J.S.S., 2011. Decoupled warming and monsoon precipitation in East Asia over the last deglaciation. *Earth Planet. Sci. Lett.* 301, 256-264.

Prins, M.A., Zheng, H., Beets, K., Troelstra, S., Bacon, P., Kamerling, I., Wester, W., Konert, M., Huang, X., Ke, W., Vandenberghe, J., 2009. Dust supply from river floodplains: the case of the lower Huang He (Yellow River) recorded in a loess-palaeosol sequence from the Mangshan Plateau. *J. Quat. Sci.* 24, 75-84.

Zheng, H., Huang, X., Ji, J., Liu, R., Zeng, Q., Jiang, F., 2007. Ultra-high rates of loess sedimentation at Zhengzhou since Stage 7: Implication for the Yellow River erosion of the Sanmen Gorge. *Geomorphology* 85, 131-142.